

**EXERCISE 8.1**

PAGE: 146

1. The angles of a quadrilateral are in the ratio 3 : 5 : 9 : 13. Find all the angles of the quadrilateral.

Solution:

Let the common ratio between the angles be  $x$ .

We know that the sum of the interior angles of the quadrilateral =  $360^\circ$

Now,

$$3x + 5x + 9x + 13x = 360^\circ$$

$$\Rightarrow 30x = 360^\circ \Rightarrow x =$$

$$12^\circ$$

, Angles of the quadrilateral are:

$$3x = 3 \times 12^\circ = 36^\circ$$


$$5x = 5 \times 12^\circ = 60^\circ$$

$$9x = 9 \times 12^\circ = 108^\circ$$

$$13x = 13 \times 12^\circ = 156^\circ$$

2. If the diagonals of a parallelogram are equal, then show that it is a rectangle.

Solution:



Given that,

$$AC = BD$$

To show that ABCD is a rectangle if the diagonals of a parallelogram are equal

To show ABCD is a rectangle, we have to prove that one of its interior angles is right-angled. Proof,



In  $\triangle ABC$  and  $\triangle BAD$ ,

$AB = BA$  (Common)

$BC = AD$  (Opposite sides of a parallelogram are equal)

$AC = BD$  (Given)

Therefore,  $\triangle ABC \cong \triangle BAD$  [SSS congruency]

$\angle A = \angle B$  [Corresponding parts of Congruent Triangles]

also,

$\angle A + \angle B = 180^\circ$  (Sum of the angles on the same side of the transversal)

$\Rightarrow 2\angle A = 180^\circ$


$\Rightarrow \angle A = 90^\circ = \angle B$

Therefore, ABCD is a rectangle.

Hence Proved.

**3. Show that if the diagonals of a quadrilateral bisect each other at right angles, then it is a rhombus.**

Solution:



Let ABCD be a quadrilateral whose diagonals bisect each other at right angles.

Given that,

$OA = OC$

$OB = OD$

and  $\angle AOB = \angle BOC = \angle OCD = \angle ODA = 90^\circ$

To show that if the diagonals of a quadrilateral bisect each other at right angles, then it is a rhombus, we have to prove that ABCD is a parallelogram and  $AB = BC = CD = AD$



Proof,

In  $\triangle AOB$  and  $\triangle COB$ ,

$OA = OC$  (Given)

$\angle AOB = \angle COB$  (Opposite sides of a parallelogram are equal)

$OB = OB$  (Common)

Therefore,  $\triangle AOB \cong \triangle COB$  [SAS congruency]

Thus,  $AB = BC$  [CPCT]

Similarly, we can prove,

$BC = CD$

$CD = AD$

$AD = AB$

,  $AB = BC = CD = AD$


Opposite sides of a quadrilateral are equal. Hence, it is a parallelogram.

ABCD is rhombus as it is a parallelogram whose diagonals intersect at a right angle.

Hence Proved.

**4. Show that the diagonals of a square are equal and bisect each other at right angles.**

Solution:



Let ABCD be a square and its diagonals AC and BD intersect each other at O.

To show that,



$$AC = BD \text{ AO} =$$

$$OC \text{ and } \angle AOB =$$

$$90^\circ$$

Proof,

In  $\triangle ABC$  and  $\triangle BAD$ ,

$$AB = BA \text{ (Common)}$$

$$\angle ABC = \angle BAD = 90^\circ$$

$$BC = AD \text{ (Given)}$$

$$\triangle ABC \cong \triangle BAD \text{ [SAS congruency]}$$

Thus,

$$AC = BD \text{ [CPCT]}$$

diagonals are equal.

Now,

In  $\triangle AOB$  and  $\triangle COD$ ,

$$\angle BAO = \angle DCO \text{ (Alternate interior angles)}$$

$$\angle AOB = \angle COD \text{ (Vertically opposite) AB}$$

$$= CD \text{ (Given)}$$

$$\therefore \triangle AOB \cong \triangle COD \text{ [AAS congruency]}$$

Thus,

$$AO = CO \text{ [CPCT].}$$

, Diagonal bisect each other

Now,

In  $\triangle AOB$  and  $\triangle COB$ ,

$$OB = OB \text{ (Given)}$$

$$AO = CO \text{ (diagonals are bisected)}$$

$$AB = CB \text{ (Sides of the square)}$$



,  $\triangle AOB \cong \triangle COB$  [SSS congruency] also,

$$\angle AOB = \angle COB$$


$$\angle AOB + \angle COB = 180^\circ \text{ (Linear pair)}$$

$$\text{Thus, } \angle AOB = \angle COB = 90^\circ$$

, Diagonals bisect each other at right angles

**5. Show that if the diagonals of a quadrilateral are equal and bisect each other at right angles, then it is a square.**

Solution:



Given that,

Let ABCD be a quadrilateral and its diagonals AC and BD bisect each other at a right angle at O.

To prove that,

The Quadrilateral ABCD is a square.

Proof,

In  $\triangle AOB$  and  $\triangle COD$ ,

$$AO = CO \text{ (Diagonals bisect each other)}$$

$$\angle AOB = \angle COD \text{ (Vertically opposite)}$$

$$OB = OD \text{ (Diagonals bisect each other)}$$

$$\text{, } \triangle AOB \cong \triangle COD \text{ [SAS congruency]}$$

Thus,

$$AB = CD \text{ [CPCT]} \text{ — (i)}$$

also,

$$\angle OAB = \angle OCD \text{ (Alternate interior angles)}$$



$\Rightarrow AB \parallel CD$

Now,

In  $\triangle AOD$  and  $\triangle COD$ ,

$AO = CO$  (Diagonals bisect each other)

$\angle AOD = \angle COD$  (Vertically opposite)

$OD = OD$  (Common)

,  $\triangle AOD \cong \triangle COD$  [SAS congruency]

Thus,

$AD = CD$  [CPCT] — (ii)

also,

$AD = BC$  and  $AD = CD$

$\Rightarrow AD = BC = CD = AB$  — (ii) also,  $\angle ADC =$

$\angle BCD$  [CPCT] and  $\angle ADC + \angle BCD = 180^\circ$

(co-interior angles)

$\Rightarrow 2\angle ADC = 180^\circ$

$\Rightarrow \angle ADC = 90^\circ$  — (iii)

One of the interior angles is a right angle.

Thus, from (i), (ii) and (iii), given quadrilateral ABCD is a square.

Hence Proved.

**6. Diagonal AC of a parallelogram ABCD bisects  $\angle A$  (see Fig. 8.19). Show that**

**(i) it bisects  $\angle C$  also,**

**(ii) ABCD is a rhombus.**




Fig. 8.19

Solution:

(i) In  $\triangle ADC$  and  $\triangle CBA$ ,

$AD = CB$  (Opposite sides of a parallelogram)

$DC = BA$  (Opposite sides of a parallelogram)

$AC = CA$  (Common Side)

$\therefore \triangle ADC \cong \triangle CBA$  [SSS congruency]

Thus,

$\angle ACD = \angle CAB$  by CPCT and

$\angle CAB = \angle CAD$  (Given)

$\Rightarrow \angle ACD = \angle BCA$

Thus,

$AC$  bisects  $\angle C$  also.

(ii)  $\angle ACD = \angle CAD$  (Proved above)

$\Rightarrow AD = CD$  (Opposite sides of equal angles of a triangle are equal)


Also,  $AB = BC = CD = DA$  (Opposite sides of a parallelogram)

Thus,

$ABCD$  is a rhombus.

**7.  $ABCD$  is a rhombus. Show that diagonal  $AC$  bisects  $\angle A$  as well as  $\angle C$  and diagonal  $BD$  bisects  $\angle B$  as well as  $\angle D$ .**

Solution:



Given that,

ABCD is a rhombus.

AC and BD are its diagonals.

Proof,

$AD = CD$  (Sides of a rhombus)

$\angle DAC = \angle DCA$  (Angles opposite of equal sides of a triangle are equal.) also,

$AB \parallel CD$

$\Rightarrow \angle DAC = \angle BCA$  (Alternate interior angles)

$\Rightarrow \angle DCA = \angle BCA$

, AC bisects  $\angle C$ .

Similarly,

We can prove that diagonal AC bisects  $\angle A$ .

Following the same method,

We can prove that the diagonal BD bisects  $\angle B$  and  $\angle D$ .


**8. ABCD is a rectangle in which diagonal AC bisects  $\angle A$  as well as  $\angle C$ . Show that:**

(i) ABCD is a square

(ii) Diagonal BD bisects  $\angle B$  as well as  $\angle D$ .

Solution:





(i)  $\angle DAC = \angle DCA$  (AC bisects  $\angle A$  as well as  $\angle C$ )

$\Rightarrow AD = CD$  (Sides opposite to equal angles of a triangle are equal) also,

$CD = AB$  (Opposite sides of a rectangle)

$AB = BC = CD = AD$

Thus, ABCD is a square.

(ii) In  $\triangle BCD$ ,

$BC = CD$

$\Rightarrow \angle CDB = \angle CBD$  (Angles opposite to equal sides are equal) also,

$\angle CDB = \angle ABD$  (Alternate interior angles)

$\Rightarrow \angle CBD = \angle ABD$

Thus, BD bisects  $\angle B$

Now,

$\angle CBD = \angle ADB$

$\Rightarrow \angle CDB = \angle ADB$

Thus, BD bisects  $\angle B$  as well as  $\angle D$ .

**9. In parallelogram ABCD, two points P and Q are taken on diagonal BD such that  $DP = BQ$  (see Fig. 8.20). Show that:**

(i)  $\triangle APD \cong \triangle CQB$

(ii)  $AP = CQ$

(iii)  $\triangle AQB \cong \triangle CPD$

(iv)  $AQ = CP$



(v) APCQ is a parallelogram




Fig. 8.20

Solution:

(i) In  $\triangle APD$  and  $\triangle CQB$ ,

$DP = BQ$  (Given)

$\angle ADP = \angle CBQ$  (Alternate interior angles)

$AD = BC$  (Opposite sides of a parallelogram)

Thus,  $\triangle APD \cong \triangle CQB$  [SAS congruency]

(ii)  $AP = CQ$  by CPCT as  $\triangle APD \cong \triangle CQB$ .

(iii) In  $\triangle AQB$  and  $\triangle CPD$ ,

$BQ = DP$  (Given)

$\angle ABQ = \angle CDP$  (Alternate interior angles)

$AB = CD$  (Opposite sides of a parallelogram)

Thus,  $\triangle AQB \cong \triangle CPD$  [SAS congruency]

(iv) As  $\triangle AQB \cong \triangle CPD$

$AQ = CP$  [CPCT]

(v) From the questions (ii) and (iv), it is clear that APCQ has equal opposite sides and also has equal and opposite angles. ,  
APCQ is a parallelogram.

10. ABCD is a parallelogram and AP and CQ are perpendiculars from vertices A and C on diagonal BD (see Fig. 8.21). Show that



(i)  $\triangle APB \cong \triangle CQD$

(ii)  $AP = CQ$




Fig. 8.21

Solution:

(i) In  $\triangle APB$  and  $\triangle CQD$ ,

$\angle ABP = \angle CDQ$  (Alternate interior angles)

$\angle APB = \angle CQD$  ( $= 90^\circ$  as  $AP$  and  $CQ$  are perpendiculars)

$AB = CD$  ( $ABCD$  is a parallelogram),

$\triangle APB \cong \triangle CQD$  [AAS congruency]

(ii) As  $\triangle APB \cong \triangle CQD$ .

,  $AP = CQ$  [CPCT]

11. In  $\triangle ABC$  and  $\triangle DEF$ ,  $AB = DE$ ,  $AB \parallel DE$ ,  $BC = EF$  and  $BC \parallel EF$ . Vertices  $A$ ,  $B$  and  $C$  are joined to vertices  $D$ ,  $E$  and  $F$ , respectively (see Fig. 8.22).

Show that

(i) quadrilateral  $ABED$  is a parallelogram

(ii) quadrilateral  $BEFC$  is a parallelogram

(iii)  $AD \parallel CF$  and  $AD = CF$

(iv) quadrilateral  $ACFD$  is a parallelogram

(v)  $AC = DF$

(vi)  $\triangle ABC \cong \triangle DEF$ .




Fig. 8.22

Solution:

(i)  $AB = DE$  and  $AB \parallel DE$  (Given)

Two opposite sides of a quadrilateral are equal and parallel to each other.

Thus, quadrilateral ABED is a parallelogram (ii)

Again  $BC = EF$  and  $BC \parallel EF$ .

Thus, quadrilateral BEFC is a parallelogram.

(iii) Since ABED and BEFC are parallelograms.

$\Rightarrow AD = BE$  and  $BE = CF$  (Opposite sides of a parallelogram are equal)

,  $AD = CF$ .

Also,  $AD \parallel BE$  and  $BE \parallel CF$  (Opposite sides of a parallelogram are parallel)

,  $AD \parallel CF$

(iv)  $AD$  and  $CF$  are opposite sides of quadrilateral ACFD which are equal and parallel to each other. Thus, it is a parallelogram.

(v) Since ACFD is a parallelogram

$AC \parallel DF$  and  $AC = DF$

(vi) In  $\triangle ABC$  and  $\triangle DEF$ ,

$AB = DE$  (Given)

$BC = EF$  (Given)

$AC = DF$  (Opposite sides of a parallelogram)

,  $\triangle ABC \cong \triangle DEF$  [SSS congruency]

12. ABCD is a trapezium in which  $AB \parallel CD$  and  $AD = BC$  (see Fig. 8.23). Show that



(i)  $\angle A = \angle B$

(ii)  $\angle C = \angle D$

(iii)  $\triangle ABC \cong \triangle BAD$

(iv) diagonal  $AC =$  diagonal  $BD$

[Hint: Extend AB and draw a line through C parallel to DA intersecting AB produced at E.]




Fig. 8.23

Solution:

To Construct: Draw a line through C parallel to DA intersecting AB produced at E.

(i)  $CE = AD$  (Opposite sides of a parallelogram)

$AD = BC$  (Given)

,  $BC = CE$

$\Rightarrow \angle CBE = \angle CEB$

also,

$\angle A + \angle CBE = 180^\circ$  (Angles on the same side of transversal and  $\angle CBE = \angle CEB$ )

$\angle B + \angle CBE = 180^\circ$  (As Linear pair)

$\Rightarrow \angle A = \angle B$

(ii)  $\angle A + \angle D = \angle B + \angle C = 180^\circ$  (Angles on the same side of transversal)

$\Rightarrow \angle A + \angle D = \angle A + \angle C$  ( $\angle A = \angle B$ )

$\Rightarrow \angle D = \angle C$

(iii) In  $\triangle ABC$  and  $\triangle BAD$ ,

$AB = AB$  (Common)

$\angle DBA = \angle CBA$



$AD = BC$  (Given)

,  $\triangle ABC \cong \triangle BAD$  [SAS congruency]

(iv) Diagonal  $AC =$  diagonal  $BD$  by CPCT as  $\triangle ABC \cong \triangle BAD$ .

EducationBharat001



## EXERCISE 8.2

PAGE: 150

1. ABCD is a quadrilateral in which P, Q, R and S are mid-points of the sides AB, BC, CD and DA (see Fig 8.29).

AC is a diagonal. Show that:

- (i)  $SR \parallel AC$  and  $SR = \frac{1}{2} AC$
- (ii)  $PQ = SR$
- (iii) PQRS is a parallelogram.




Fig. 8.29

Solution:

(i) In  $\triangle DAC$ ,

R is the mid point of DC and S is the mid point of DA.

Thus by mid point theorem,  $SR \parallel AC$  and  $SR = \frac{1}{2} AC$

(ii) In  $\triangle BAC$ ,

P is the mid point of AB and Q is the mid point of BC.

Thus by mid point theorem,  $PQ \parallel AC$  and  $PQ = \frac{1}{2} AC$  also,

$$SR = \frac{1}{2} AC$$

$$, PQ = SR$$

(iii)  $SR \parallel AC$  from question (i) and,

$PQ \parallel AC$  from question (ii)

$\Rightarrow SR \parallel PQ$  from (i) and (ii) also,


$$PQ = SR$$

, PQRS is a parallelogram.

2. ABCD is a rhombus and P, Q, R and S are the mid-points of the sides AB, BC, CD and DA, respectively. Show that the quadrilateral PQRS is a rectangle.



Solution:



Given in the question,

ABCD is a rhombus and P, Q, R and S are the mid-points of the sides AB, BC, CD and DA, respectively. To

Prove,

PQRS is a rectangle.

Construction, Join

AC and BD.

Proof:

In  $\triangle DRS$  and  $\triangle BPQ$ ,

$DS = BQ$  (Halves of the opposite sides of the rhombus)

$\angle SDR = \angle QBP$  (Opposite angles of the rhombus)

$DR = BP$  (Halves of the opposite sides of the rhombus)

$\therefore \triangle DRS \cong \triangle BPQ$  [SAS congruency]

$RS = PQ$  [CPCT] ————— (i)

In  $\triangle QCR$  and  $\triangle SAP$ ,

$RC = PA$  (Halves of the opposite sides of the rhombus)

$\angle RCQ = \angle PAS$  (Opposite angles of the rhombus)

$CQ = AS$  (Halves of the opposite sides of the rhombus)





,  $\triangle QCR \cong \triangle SAP$  [SAS congruency]

$RQ = SP$  [CPCT]————— (ii)

Now,

In  $\triangle CDB$ ,

R and Q are the mid points of CD and BC, respectively.

$\Rightarrow QR \parallel BD$

also,

P and S are the mid points of AD and AB, respectively.

$\Rightarrow PS \parallel BD$

$\Rightarrow QR \parallel PS$

, PQRS is a parallelogram.

also,  $\angle PQR = 90^\circ$

Now,

In PQRS,


$RS = PQ$  and  $RQ = SP$  from (i) and (ii)

$\angle Q = 90^\circ$

, PQRS is a rectangle.

**3. ABCD is a rectangle and P, Q, R and S are mid-points of the sides AB, BC, CD and DA, respectively. Show that the quadrilateral PQRS is a rhombus.**

Solution:



Given in the question,



ABCD is a rectangle and P, Q, R and S are mid-points of the sides AB, BC, CD and DA, respectively. Construction,

Join AC and BD.

To Prove, PQRS is

a rhombus.

Proof:

In  $\triangle ABC$

P and Q are the mid-points of AB and BC, respectively

,  $PQ \parallel AC$  and  $PQ = \frac{1}{2} AC$  (Midpoint theorem) — (i)

In  $\triangle ADC$ ,

$SR \parallel AC$  and  $SR = \frac{1}{2} AC$  (Midpoint theorem) — (ii)

So,  $PQ \parallel SR$  and  $PQ = SR$

As in quadrilateral PQRS one pair of opposite sides is equal and parallel to each other, so, it is a parallelogram.

,  $PS \parallel QR$  and  $PS = QR$  (Opposite sides of parallelogram) — (iii)

Now,

In  $\triangle BCD$ ,

Q and R are mid points of side BC and CD, respectively.

,  $QR \parallel BD$  and  $QR = \frac{1}{2} BD$  (Midpoint theorem) — (iv)

$AC = BD$  (Diagonals of a rectangle are equal) — (v)

From equations (i), (ii), (iii), (iv) and (v),

$PQ = QR = SR = PS$

So, PQRS is a rhombus.

Hence Proved

4. ABCD is a trapezium in which  $AB \parallel DC$ , BD is a diagonal and E is the mid-point of AD. A line is drawn through E parallel to AB intersecting BC at F (see Fig. 8.30). Show that F is the mid-point of BC.




Fig. 8.30

Solution:

Given that,

ABCD is a trapezium in which  $AB \parallel DC$ , BD is a diagonal and E is the mid-point of AD.

To prove,

F is the mid-point of BC.

Proof,

BD intersected EF at G.

In  $\triangle BAD$ ,

E is the mid point of AD and also  $EG \parallel AB$ .

Thus, G is the mid point of BD (Converse of mid point theorem)


Now,

In  $\triangle BDC$ ,

G is the mid point of BD and also  $GF \parallel AB \parallel DC$ .

Thus, F is the mid point of BC (Converse of mid point theorem)

**5. In a parallelogram ABCD, E and F are the mid-points of sides AB and CD, respectively (see Fig. 8.31). Show that the line segments AF and EC trisect the diagonal BD.**



Solution:

Given that,

ABCD is a parallelogram. E and F are the mid-points of sides AB and CD, respectively.

To show,

AF and EC trisect the diagonal BD.

Proof,

ABCD is a parallelogram

,  $AB \parallel CD$  also,

$AE \parallel FC$

Now,

$AB = CD$  (Opposite sides of parallelogram ABCD)  $\Rightarrow$

$\frac{1}{2} AB = \frac{1}{2} CD$

$\Rightarrow AE = FC$  (E and F are midpoints of side AB and CD)

AECF is a parallelogram (AE and CF are parallel and equal to each other)

$AF \parallel EC$  (Opposite sides of a parallelogram)

Now,

In  $\triangle DQC$ ,

F is mid point of side DC and  $FP \parallel CQ$  (as  $AF \parallel EC$ ).

P is the mid-point of DQ (Converse of mid-point theorem)

$\Rightarrow DP = PQ$  — (i)



Similarly,

In  $\triangle APB$ ,

E is midpoint of side AB and  $EQ \parallel AP$  (as  $AF \parallel EC$ ).

Q is the mid-point of PB (Converse of mid-point theorem)

$$\Rightarrow PQ = QB \text{ — (ii)}$$

From equations (i) and (ii),


$$DP = PQ = BQ$$

Hence, the line segments AF and EC trisect the diagonal BD.

Hence Proved.

**6. Show that the line segments joining the mid-points of the opposite sides of a quadrilateral bisect each other.**

Solution:



Let ABCD be a quadrilateral and P, Q, R and S the mid points of AB, BC, CD and DA, respectively.

Now,

In  $\triangle ACD$ ,

R and S are the mid points of CD and DA, respectively.

$$, SR \parallel AC$$

Similarly we can show that,

$$PQ \parallel AC,$$

$$PS \parallel BD \text{ and}$$

$$QR \parallel BD$$




, PQRS is parallelogram.

PR and QS are the diagonals of the parallelogram PQRS. So, they will bisect each other.

7. ABC is a triangle right angled at C. A line through the mid-point M of hypotenuse AB and parallel to BC intersects AC at D. Show that

(i) D is the mid-point of AC

(ii)  $MD \perp AC$  (iii)  $CM = MA = \frac{1}{2} AB$  Solution:



(i) In  $\triangle ACB$ ,

M is the midpoint of AB and  $MD \parallel BC$

, D is the midpoint of AC (Converse of mid point theorem)

(ii)  $\angle ACB = \angle ADM$  (Corresponding angles) also,  $\angle ACB = 90^\circ$

,  $\angle ADM = 90^\circ$  and  $MD \perp AC$

(iii) In  $\triangle AMD$  and  $\triangle CMD$ ,

$AD = CD$  (D is the midpoint of side AC)

$\angle ADM = \angle CDM$  (Each  $90^\circ$ )

$DM = DM$  (common)

,  $\triangle AMD \cong \triangle CMD$  [SAS congruency]

$AM = CM$  [CPCT] also,  $AM = \frac{1}{2} AB$  (M

is midpoint of AB)

Hence,  $CM = MA = \frac{1}{2} AB$



EducationBharat001

NCERT Solutions for Class 9 Maths Chapter 8 –  
Quadrilaterals

EducationBharat001

<https://educationbharat001.com/>